



Mark Scheme (Results)

Summer 2025

Pearson Edexcel GCSE
In Combined Science Chemistry
(1SC0) Paper 1CH

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2025

Question Paper Log Number P75513RA

Publications Code 1SC0_1CH_2506_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Combined Science- Chemistry 1SC0_1CH

Question number	Answer	Additional guidance	Mark									
1(a)(i)	<table><tr><td>proton</td><td>+1 (1)</td><td>1</td></tr><tr><td>neutron</td><td>0</td><td>1 (1)</td></tr><tr><td>electron</td><td>-1 (1)</td><td>negligible</td></tr></table>	proton	+1 (1)	1	neutron	0	1 (1)	electron	-1 (1)	negligible	allow 1(+) / +, reject -1 reject +1, -1 allow (1)-, reject +1 / 1	(3) A01-1
proton	+1 (1)	1										
neutron	0	1 (1)										
electron	-1 (1)	negligible										

Question number	Answer	Additional guidance	Mark
1(a)(ii)	A description including 10 protons (1) 10 electrons (1)	allow {'how many'/'number of'} protons allow {'how many'/'number of'} electrons / (electronic configuration) 2.8 allow 'number of protons and (number of) electrons' (2) ignore any reference to location of subatomic particles ignore neutrons	(2) A01-1

Question number	Answer	Additional guidance	Mark
1(a)(iii)	An explanation linking <u>same number of</u> protons and electrons (1) (so) the (+ and -) <u>charges</u> cancel (1)	ignore reference to neutrons allow <u>charge</u> (on neon atom) = <u>0</u> +10 plus -10 = 0 (2)	(2) A02-1

Question number	Answer	Additional guidance	Mark
1(b)	$90.48 \times 20 + 0.27 \times 21 + 9.25 \times 22 = 2018.77 \text{ (1)}$ $\frac{2018.77}{100} = 20.1877 \text{ (1)}$	20.2 / 20.19 with or without working scores 2 marks 20.18 / 20.187 with or without working scores 1 mark 2018.77 with or without working (1) allow $90.48 \times 20 + 0.27 \times 21 + 9.25 \times 22$ unevaluated (1) or $90.48 \times 0.20 + 0.27 \times 0.21 + 9.25 \times 0.22$ unevaluated (1) incorrect / early rounding loses MP2 allow 3 or more sf	(2) A02-1

Total for Question 1 = 9 marks

Question number	Answer	Additional guidance	Mark
2(a)(i)	aq (1) KNO ₃ (1)	allow AQ / Aq / aQ reject aqueous allow NO ₃ K / K(NO ₃) / (K)NO ₃ reject any other answer all letters must be capitals. The 3 must be a subscript. ignore any brackets in formula	(2) A02-1

Question number	Answer	Additional guidance	Mark
2(a)(ii)	silver chloride	reject silver chlorine ignore solid / precipitate / ppt	(1) A02-1

Question number	Answer	Additional guidance	Mark
2(a)(iii)	An explanation including any 3 from - filtration (1) <u>pour mixture through</u> {filter funnel / filter paper} (1) {precipitate / residue / solid / AgCl} stays on the paper (1) {solution / filtrate / liquid} goes through (the paper) (1)	allow to filter (mixture) allow <u>pour mixture through</u> {filter / funnel} allow precipitate is left behind (in filter / funnel) allow particles too big to go through ignore mixture goes through ignore any additional processes after filtration step if filtration occurs after any other process, e.g. heating, MP1 and MP2 do not score but MP3 and MP4 can be scored diagram can score all MPs	(3) A03-3

Question number	Answer	Additional guidance	Mark
2(b)	A description including • {silver nitrate / solution} has ions (1) • {silver / solid / metal} has delocalised electrons (1) • {ions / electrons} move (1)	reject silver nitrate has electrons {that move / delocalised} for MP1 allow {free / sea of} electrons reject <u>silver</u> has ions {that move / delocalised} for MP2 allow charged particles move MP3 for idea of movement of charge (even if, for example, MP1 and/or MP2 reversed or wrong)	(3) A02-1

Total for Question 2 = 9 marks

Question number	Answer	Additional guidance	Mark
3(a)	An explanation linking - period 2 (1) (because) 2 (electron) shells (1)	accept 'row', <u>horizontal</u> line allow electron configurations for each element ignore same number of shells reject 2 outer shells if no other mark awarded period = number of electron shells (1)	(2) AO1.1


Question number	Answer	Additional guidance	Mark
3(b)(i)	An explanation linking - simple covalent / simple molecular (1) (because) <u>weak</u> intermolecular forces (of attraction) (1) (so) only a small / low amount of {energy/heat} is needed (to break these forces) (1)	reject ionic / ions for MP1 & MP2 only ignore covalent alone allow weak intermolecular bonds reject weak covalent bonds reject hydrogen bonds reject weak (intermolecular) forces between bonds ignore less {energy / heat} reject breaking {covalent / ionic} bonds	(3) AO1.1

Question number	Answer	Mark
3(b)(ii)	C less less is the only correct answer A is incorrect as the particles will have less energy B is incorrect as the particles will move less D is incorrect as the particles will have less energy and move less	(1) AO1.1

Question number	Answer	Mark
3(b)(iii)	B Fractional distillation A, C and D are not correct as they would not separate oxygen and nitrogen	(1) A02.2

Question number	Answer	Additional guidance	Mark
3(c)	$\text{N}_2 + 2\text{O}_2 \rightarrow 2\text{NO}_2$ (2) all formulae correct (1) balancing of <u>correct formulae</u> for the reaction (1)	do not penalise incorrect cases accept = / $\rightleftharpoons$ ignore state symbols reject superscripts allow multiples	(2) A02.1

Total for Question 3 = 9 marks

Question number	Answer	Mark
4(a)(i)	 C B and D are not correct as the carbon dioxide would escape A is not correct as the bung would pop out and the carbon dioxide would escape	(1) A03.3

Question number	Answer	Additional guidance	Mark
4(a)(ii)	calcium carbonate	allow CaCO ₃	(1) A02.2

Question number	Answer	Additional guidance	Mark
4(a)(iii)	formula mass of Cu₂CO₃(OH)₂: (63.5×2)+12+(16×3)+((16+1)×2) = 221 (1) $\frac{22.1}{221} = 0.1$ (1) 0.1 × 2 = 0.2 (1) 0.2 × 79.5 = 15.9 (1)	15.9 gains 4 marks with or without working allow ECF throughout alternative method: (63.5×2)+12+(16×3)+((16+1)×2) = 221 (1) mole ratio 1 mole: 2 moles CuO (1) 221g malachite : 2 × 79.5 = 159g CuO (1) 22.1 malachite : 159 × 22.1/221 = 15.9g (1) 31.8 (g) scores 3 7.95 (g) scores 3	(4) A02.1

Question number	Answer	Additional guidance	Mark
4(b)	comments to include three from: method 1 - mixing - strength - means more will react OR method 2 - not mixing - weakness - less will react (1) method 1 and / or method 2 - quantity of carbon not given – weakness - may not be in excess (1) method 1 - use of a metal crucible - strength - conducts more heat (1) method 1 - use of a metal crucible – weakness – metal crucible might react (1) method 1 – transfers reactants between containers – weakness – loss of reactants (1) method 1 and / or method 2 - roaring flame - strength - gives enough energy for reaction to occur (1) method 1 and / or method 2 – separation of copper – weakness - no method given (1)	Note: implication of strength/weakness and method can be obtained from the way in which the candidate gives the answer eg using the metal crucible is better as it conducts more heat Only Method 1 is using the metal crucible so the method has been identified. 'is better' would be another way of saying that it is a strength	(3) A03.1

Total for Question 4= 9 marks

Question number	Answer	Mark
5(a)	D sodium sulfate A, B and C are incorrect as they are not soluble in water	(1) A01.2 comp

Question number	Answer	Additional guidance	Mark
5(b)(i)	$\frac{64.0}{120} = 0.533333$ (1) $0.533333 \times 100 = 53.333$ (1) (%) 53.3 (1)	53.3 with or without working gains 3 marks 53(.33)% with or without working gains 2 marks allow ecf for MP3 only if appropriate Ar values have been used and answer <100% ignore any dot above the 3 13.3 scores 2 marks $\frac{16.0}{120} \times 100$ unevaluated scores 1	(3) A02.1

Question number	Answer	Additional guidance	Mark
5(b)(ii)	$\frac{7.20}{120} = 0.06$ (1) $0.06 \times 4 = 0.24$ (1) $0.24 \times 6.02 \times 10^{23} = 1.444(8) \times 10^{23}$ (1)	$1.444(8) \times 10^{23}$ with or without working gains 3 marks Allow any number of sig figs except 1 allow ECF throughout MP3 only awarded ECF if numbers from the question are multiplied by <i>L</i> $120/7.2 = 16.67$ (0) $16.67 \times 4 = 66.67$ (1) $66.67 \times L = 4.01 \times 10^{25}$ (1) ie scores 2 (missed mp1) $0.06 \times L = 3.612 \times 10^{22}$ scores 2 (missed mp2)	(3) A02.1

Question number	Indicative content	Mark
5(c)	Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlined in the generic mark scheme. The indicative content below is not prescriptive and candidates are not required to include all the material that is indicated as relevant. Additional content included in the response must be scientific and relevant. AO1 (6 marks) • Mg^{2+} and SO_4^{2-} ions are present in magnesium sulfate solution • water splits to form H^+ and OH^- ions • negative ions / anions move positive electrode / anode • sulfate ions / SO_4^{2-} and hydroxide / OH^- ions move to anode • can credit ions shown on diagram • hydroxide ions / OH^- lose electrons • loss of electrons is oxidation • hydroxide ions are oxidised • $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^-$ • positive ions / cations move to negative electrode / cathode • magnesium ions / Mg^{2+} and hydrogen ions / H^+ move to cathode • hydrogen forms as it's less reactive than magnesium • hydrogen ions / H^+ gain electrons • gain of electrons is reduction • hydrogen ions are reduced • $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$	(6) AO1.1

Level	Mark	Descriptor
	0	• No rewardable material.
Level 1	1-2	• Demonstrates elements of chemical understanding, some of which is inaccurate. Understanding of scientific ideas lacks detail. (AO1) • Presents an explanation with some structure and coherence. (AO1)
Level 2	3-4	• Demonstrates chemical understanding, which is mostly relevant but may include some inaccuracies. Understanding of scientific ideas is not fully detailed and/or developed. (AO1) • Presents an explanation that has a structure which is mostly clear, coherent and logical. (AO1)
Level 3	5-6	• Demonstrates accurate and relevant chemical understanding throughout. Understanding of the scientific ideas is detailed and fully developed. (AO1) • Presents an explanation that has a well-developed structure which is clear, coherent and logical. (AO1)

Level	Mark	Descriptor	Additional Guidance
	0	No rewardable material.	Read whole answer and ignore all incorrect material and any references to other metals / discard any contradictory material then:
Level 1	1–2	<u>Additional Guidance</u> Identifies ions from magnesium sulfate Identifies ions from water Identifies correct movement of ions Oxidation & reduction	<u>Possible candidate response</u> (all examples, not a definitive list) Mg^{2+} / SO_4^{2-} / H^+ / OH^- ions (any one ion) present (1) Identifies at least two ions present (2) Negative ions move to anode (1) Negative ions move to {anode/positive electrode} and positive ions move to {cathode/negative electrode} (2) Oxidation at anode and reduction at cathode (2)
Level 2	3–4	<u>Additional Guidance</u> Identifies one ion and states where it moves to during electrolysis Identifies at least 2 ions present and their movement during electrolysis Identifies H^+ or OH^- , states direction of movement and product formed Describes oxidation and reduction	<u>Possible candidate response</u> Mg^{2+} present and moves to cathode (3) Mg^{2+} and SO_4^{2-} ions are present, Mg^{2+} moves to negative electrode, SO_4^{2-} moves to the positive electrode (4) Hydrogen ions present, move to negative electrode / cathode and form hydrogen (4) Oxidation is loss of electrons at anode; reduction is gain of electrons at cathode (4)
Level 3	5–6	<u>Additional Guidance</u> Identifies both H^+ and OH^- , states direction of movement and products formed Identifies all ions present and gives the direction of movement to correct electrodes Identifies at least one cation and at least one anion, states direction of movement of ions, what happens at electrode (loss/gain of electrons) for one ion and identifies product Identifies at least one cation and at least one anion, states direction of movement of both ions, what happens at electrode (loss/gain of electrons or reduction/oxidation) for both ions and identifies products; MAY attempt half equations, but not essential Correctly write both half equations	<u>Possible candidate response</u> Hydrogen ions and hydroxide ions present, H^+ move to negative electrode form hydrogen, OH^- move to anode and form oxygen (5) Magnesium, sulfate, hydrogen and hydroxide ions are present, the magnesium and hydrogen ions move towards the negative electrode, sulfate and hydroxide move towards the positive electrode (5) Mg^{2+} ions are present; H^+ move to cathode, the H^+ gain electrons OR are reduced to form H_2 . (5) Mg^{2+} and SO_4^{2-} are present from the magnesium sulfate, H^+ and OH^- are present from the water. Mg^{2+} and H^+ move to cathode, the H^+ gains electrons and is reduced to form H_2 . (6) Mg^{2+} and SO_4^{2-} are present from the sodium sulfate, H^+ and OH^- are present from the water. Na^+ and H^+ move to cathode, the hydrogen is reduced to form H_2 . OH^- and SO_4^{2-} move to anode and oxygen is produced; $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^-$ (6) $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ and $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^-$ (6)

Total for Question 5 = 13 marks

Question number	Answer	Additional guidance	Mark
6(a)	An answer linking any two from - so that the ammonia can {react with / dissolve in} the water (1) (so) colour change seen (1) - hydroxide ions form (1)	allow 'dampness' for water allow ammonia is absorbed by water ignore ammonia condenses on / sticks to paper ignore any colour even if incorrect but reject bleach for MP2 allow alkaline / basic solution formed	(2) A02.2

Question number	Answer	Additional guidance	Mark
6(b)(i)	An explanation linking - both reactions happen at the same rate (1) (because) there is no overall change in the system (1)	ignore both reactions happen at the same time allow total amount/concentration of products and reactants are <u>constant</u> ignore reference to amounts/concentration of products and reactants being <u>equal</u> ignore closed system	(2) A01.1

Question number	Answer	Mark
6(b)(ii)	A 200 1000 B , C and D are incorrect as a temperature of 200°C and a pressure of 1000 would give the highest percentage yield of ammonia	(1) A03.1

Question number	Answer	Additional guidance	Mark
6(b)(iii)	An explanation linking: - equilibrium would move {backwards / to left} (1) (because) there are more moles on {left hand side / reactants} (1)	allow references to lower yield / less product formed / more reactant formed. allow particles / molecules for moles allow RA throughout if no other mark scored, 'equilibrium shifts to the side with more moles / molecules' scores 1 mark. ignore all references to temperature / heat / exothermic / endothermic. ignore all references to rate.	(2) A01.1

Question number	Answer	Additional guidance	Mark
6(c)	An explanation linking: - ammonia and hydrogen chloride / gases form (in the forward reaction) (1) - the gas / gases rise to the top of the container (1) - where it is cooler / the gas(es) cool (1) - the reverse reaction takes place / reaction forms {white solid/ammonium chloride} (at the top of tube) (1)	allow formula throughout allow endothermic reaction takes place reject ammonium chloride gas / hydrochloric acid forms for MP1 only allow exothermic reaction takes place allow hydrogen chloride and ammonia react (at the top of the tube)	(4) A03.2

Total for Question 6 = 11 marks

