

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

9701/24

May/June 2026

1 hour 15 minutes

No additional materials are needed.

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

- 1 Aluminium is a Group 13 element that forms a number of compounds.

Table 1.1

atom or ion	number of protons	number of neutrons
^{27}Al		
$^{28}\text{Al}^{3+}$		

- (a) (i) Complete Table 1.1 to show the number of particles in each atom or ion. [2]
- (ii) Complete Figure 1.1 to show the electronic configuration of an aluminium atom.




Figure 1.1

- (b) Deduce the formula of aluminium nitrate(V).

..... [1]

- (c) AlCl_3 dimerises to form Al_2Cl_6 . Al_2Cl_6 is shown in Figure 1.2.




Figure 1.2

Describe how coordinate (dative covalent) bonds form when two AlCl_3 molecules combine to form Al_2Cl_6 .

.....

..... [1]

- (d) Aluminium forms the AlCl_4^- ion. The dot-and-cross diagram for AlCl_4^- in Figure 1.3 shows the arrangement of outer electrons.




Figure 1.3

- (i) Use Figure 1.3 and VSEPR theory to name the shape of the AlCl_4^- ion and to predict the $\text{Cl}-\text{Al}-\text{Cl}$ bond angle.

name of shape

bond angle

[2]

- (ii) Draw the three-dimensional shape of the AlCl_4^- ion.

[1]

[Total: 8]

- 2 (a) (i) Nitrogen, N_2 , contains sigma (σ) and pi (π) bonds.

Describe how a pi (π) bond is formed.

.....

 [2]

- (ii) Explain the lack of reactivity of nitrogen.

.....

 [2]

- (b) Nitrogen is used to manufacture ammonia, NH_3 , in the Haber process. An iron catalyst is used.

- (i) Complete the reaction pathway diagram in Figure 2.1 to show the effect of adding an iron catalyst to the reaction of nitrogen and hydrogen.




Figure 2.1

[1]


- (ii) State the temperature and pressure used in the Haber process.

temperature =

pressure =

[2]

- (c) The equation for the Haber process is shown.



Three separate experiments are carried out using mixtures of $\text{N}_2(\text{g})$ and $\text{H}_2(\text{g})$ to find the amount of $\text{NH}_3(\text{g})$ produced at equilibria under different conditions.

- (i) In experiment 1, 3.55 mol of $\text{N}_2(\text{g})$ and 8.60 mol of $\text{H}_2(\text{g})$ are mixed and left to reach equilibrium.

At equilibrium, 3.24 mol of NH_3 is present.

Complete Table 2.1.

Table 2.1

	$\text{N}_2(\text{g})$	$\text{H}_2(\text{g})$	$\text{NH}_3(\text{g})$
initial amount/mol	3.55	8.60	0
change in amount/mol			3.24
amount at equilibrium/mol			3.24

[3]

- (ii) In experiment 2, under different conditions $\text{N}_2(\text{g})$ and $\text{H}_2(\text{g})$ are mixed and left to reach equilibrium.

Table 2.2 shows the amount, in mol, at equilibrium of $\text{N}_2(\text{g})$, $\text{H}_2(\text{g})$ and $\text{NH}_3(\text{g})$.

Table 2.2

	$\text{N}_2(\text{g})$	$\text{H}_2(\text{g})$	$\text{NH}_3(\text{g})$
amount at equilibrium/mol	0.874	1.340	0.347

Calculate the mole fraction of $\text{NH}_3(\text{g})$ in the equilibrium mixture in experiment 2.

mole fraction of $\text{NH}_3(\text{g})$ = [2]

- (iii) The expression for the equilibrium constant, K_p , for the Haber process is shown.

$$K_p = \frac{p(\text{NH}_3)^2}{p(\text{N}_2) \times p(\text{H}_2)^3}$$

Describe what is meant by partial pressure.

.....

 [1]

- (iv) In experiment 3, the mixture at equilibrium contains the following partial pressures of $\text{N}_2(\text{g})$ and $\text{H}_2(\text{g})$.

$$p(\text{N}_2) = 1590 \text{ kPa}$$

$$p(\text{H}_2) = 1750 \text{ kPa}$$

For experiment 3, the equilibrium constant, K_p , is $8.49 \times 10^{-7} \text{ kPa}^{-2}$.

Calculate the partial pressure, in kPa, of $\text{NH}_3(\text{g})$ at equilibrium.

$$p(\text{NH}_3) = \dots\dots\dots \text{ kPa} \quad [3]$$

(d) Explain the effect of increasing temperature on rate of reaction.

.....

.....

.....

..... [3]

(e) Figure 2.2 shows the Boltzmann distribution for a sample of X(g) at temperature T_1 .




Figure 2.2

Sketch a curve on Figure 2.2 to show the Boltzmann distribution for the sample of X(g) at a temperature higher than T_1 . [2]

[Total: 21]

- 3 (a) (i) State the trend in atomic radius of the elements across Period 3.

..... [1]

- (ii) State the formula of the oxide formed when sodium reacts with oxygen.

..... [1]

- (iii) Write an equation for the reaction of MgO with water.

..... [1]

- (b) Table 3.1 shows some of the physical properties of Period 3 chlorides.

Table 3.1

	melting point /°C	boiling point /°C	type of bonding	type of structure
NaCl	801	1413		
SiCl₄	–69	58		

- (i) Complete Table 3.1 using the abbreviations shown.

M = metallic
C = covalent
I = ionic
S = simple
G = giant

[2]

- (ii) State the formula of the phosphorus-containing product formed when PCl_5 reacts with an excess of water. Suggest the pH of the resulting solution.

formula =

pH of solution =

[2]

- (c) (i) Describe what is observed when white solid strontium nitrate decomposes on heating.

..... [1]

- (ii) State the trend in solubility down the Group 2 hydroxides.

..... [1]

[Total: 9]

- 4 (a) Alcohol **S** has the structural formula $\text{CH}_3\text{C}(\text{OH})(\text{CH}_3)\text{CH}_2\text{CH}_3$.

Classify alcohol **S**.

..... [1]

- (b) Figure 4.1 shows compound **T**.




Figure 4.1

Tests are carried out on **T** using different reagents.

Complete Table 4.1 to show the observations of each test on **T**. If no reaction occurs, write 'no reaction' in the relevant box.

Table 4.1

reagent	observations
Na	
acidified $\text{K}_2\text{Cr}_2\text{O}_7$	
alkaline $\text{I}_2(\text{aq})$	

[3]

(c) Compound **W** is an aliphatic primary alcohol.

W is oxidised by acidified KMnO_4 when heated under reflux. Figure 4.2 shows the infrared spectrum of the resulting product.




Figure 4.2

Table 4.2

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/ cm^{-1}
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3650

The infrared spectrum in Figure 4.2 confirms that **W** is a primary alcohol.

Use Figure 4.2 to identify **two** absorbances that confirm **W** is a primary alcohol. State the bond and functional group containing the bond for each absorbance.

.....

 [2]

(d) Butane is converted to pentanenitrile via a two-step synthesis, as shown in Figure 4.3.




Figure 4.3

(i) Suggest the reagent and conditions for each step in Figure 4.3.

step 1
 step 2 [2]

(ii) Pentanenitrile reacts with dilute acid.

Draw the displayed formula of the organic product made in this reaction.

[1]

[Total: 9]

5 Long-chain alkanes from crude oil can be cracked to make alkenes.

(a) State the conditions used for the cracking of alkanes.

.....
..... [1]

(b) Propene is an alkene that is made by cracking. Alkenes are used in addition polymerisation.

(i) Draw **one** repeat unit of the addition polymer formed from propene.

[1]

(ii) The disposal of poly(alkenes) by combustion produces harmful products.

State **one** other disadvantage associated with the disposal of poly(alkenes) such as poly(propene).

.....
..... [1]

(iii) Aqueous bromine is used to test for the presence of the C=C bond in propene.

State what is observed when aqueous bromine is mixed with propene.

..... [1]

(c) But-1-ene and but-2-ene are structural isomers.

State the type of structural isomerism shown by but-1-ene and but-2-ene.

..... [1]

(d) But-2-ene reacts with HBr(g) . The HBr acts as an electrophile in the reaction.

(i) State how HBr acts as an electrophile in terms of electron transfer.

..... [1]

(ii) Complete Figure 5.1 to show the mechanism for the reaction of but-2-ene with HBr .

Include charges, dipoles, lone pairs of electrons and curly arrows, as appropriate.



Figure 5.1

[4]

(e) But-1-ene and bromine, Br_2 , react in an electrophilic addition reaction to make product **P**.

Give the systematic name for **P**.

..... [1]

- (f) Figure 5.2 shows the structure of methylpropene.




Figure 5.2

Methylpropene reacts with HCl to produce products **Q** and **R**.




Figure 5.3

Explain why **R** is the major product in this reaction.

.....

.....

.....

..... [2]

[Total: 13]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

The Periodic Table of Elements

Group																							
1	2													13	14	15	16	17	18				
														1 H hydrogen 1.0					2 He helium 4.0				
3 Li lithium 6.9	4 Be beryllium 9.0	Key atomic number atomic symbol name relative atomic mass																5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
	11 Na sodium 23.0	12 Mg magnesium 24.3																	13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8						
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3						
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —					
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —					

lanthanoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

actinoids